

Products to Discontinue

Important: Please discontinue blood thinners two weeks before any scheduled procedure. Aspirin, arthritis medications, cold medicines, and other drugs containing aspirin or anti-inflammatories can cause excessive bleeding during surgical procedures. The list below is not a complete list of aspirin-containing or otherwise prohibited products, so be sure to check all medications you might take in the two weeks prior to your procedure to see if they contain aspirin. If you're not sure, contact your pharmacist.

NOTE: Do not take the following for two weeks prior to your procedure without checking with our office. If you have any questions, or are medically required to take blood thinners such as **Coumadin (warfarin), Eliquis, Xarelto, Pradaxa, Savaysa, Plavix, or Heparin**, *please notify us **AND** confirm with your doctor that it is okay to stop in anticipation of your surgical procedure.*

Advil	Enteric	Motrin
Aggrenox	Ephedra	Naprosyn
Aleve	Excedrin	Norgesic
Alka Seltzer	Fiorina	Nuprin
Anacin	Fish Oil	Pentasa
Arthritis Pain Formula	4-Way Cold Tablets	Pepto Bismol
Ascription	Garlic	Plavix
Aspirin	Ginger	Presistin
Bayer	Gingko Biloba	Propoxyphene
Bufferin	Ginseng	Saw Palmetto
Cama arthritis reliever	Glucosamine	St. John's Wort
Celebrex	Goldenseal	Trigestic
Chondroitin	Indocin	Valerian
Coricidin	Kava	Vanquish
Doans Pain Pills	Milk Thistle	Vicoprofen
Echinacea	Midol	Vitamin E
Ecotrin Emprin Compound	Mobic	

Signature: _____

Date: _____